



Search PubMed for

Go

Clear

Limits

Preview/Index

History

Clipboard

Details

Display

Abstract

Sort

Save

Text

Clip Add

Order

Entrez PubMed

PubMed Services

Related Resources

1: Eur J Immunol 1995 Feb;25(2):399-404

Related Articles, Books

Reconstitution of a functional interleukin (IL)-7 receptor demonstrates that the IL-2 receptor gamma chain is required for IL-7 signal transduction.**Ziegler SE, Morella KK, Anderson D, Kumaki N, Leonard WJ, Cosman D, Baumann H.**

Department of Molecular Biology, Immunex Research and Development Corp., Seattle.

The interleukin (IL)-2 receptor gamma chain has recently been shown to be a component of the IL-7 and IL-4 receptors. Using a transient transfection assay and the trans-activation of reporter gene constructs which are under the control of cytokine-responsive promoter elements, we have studied signal transduction through the IL-7 receptor (IL-7R). The reporter gene expression was not stimulated by receptors that contained the cytoplasmic domain of the IL-7R, either as intact IL-7R or as part of a chimeric receptor. However, co-expression of the IL-7R with the IL-2 receptor gamma chain was able to stimulate gene activation. For maximal stimulation the intact cytoplasmic domains of each chain was required.

PMID: 7875201 [PubMed - indexed for MEDLINE]

Display

Abstract

Sort

Save

Text

Clip Add

Order

[Write to the Help Desk](#)[NCBI | NLM | NIH](#)[Department of Health & Human Services](#)[Freedom of Information Act | Disclaimer](#)

Source: PubMed, 2001 Dec 19 PM 19921